Shaped excrements can be organized as single units which either are loose or dense packed or form more or less coalescing units. Excrements of all shapes and organizations can partly or completely fill faunal voids. Mites, enchytraeids, diplopods and isopods generally produce accumulations of individual units, while earthworm excreta is more or less coalesced. Depending on the species, faunal voids can also be infilled with shapeless excreta (e.g. earthworm species), or be infilled with soil material by the soil fauna (e.g. dung beetle).

The impact of soil fauna is not only related to their population size but also to their mobility and activity. Sessile soil fauna, even if numerous, may have less impact on the soil architecture than mobile species, which always move around in search for food, protection or oviposition sites. On the other hand, a few distinct faunal channels produced by sessile soil fauna can have far more impact on soil physical parameters (Edwards et al., 1993) than the partly filled in tracks with soil material and/or excrements of the mobile soil fauna. In temperate climates and in forests the impact of the soil fauna is highest in the A horizon. In climates with extremes in temperature and dryness the highest impact is found in deeper horizons (Kooistra, 1982). Faunal voids often cross or follow for some distance cracks or root channels. In arable land soil fauna can produce more voids systems than plant roots, which mainly follow existing voids.

4. Human Impact

Human impact on soil architecture can be due to soil tillage, traffic (esp. at harvest time) and slaking. Soil tillage practices, especially the main or primary tillage, have the largest influence. During tillage the soil is loosened and partly pulverized as well as often mixed and inverted, whereby forming aggregates separated by rough-walled, irregular voids. Afterwards, by natural settlement, traffic and other cultivation activities, the aggregates merge. Thus, the continuity of the tillage-induced voids decrease, while isolated voids remain (see Figure 12). In the first month after ploughing and during the seedbed preparation most of the increased porosity disappears (by about 40% at both times) (Kuipers and Van Ouwerkerk, 1963). The soil is again loosened after the harvest. Tillage voids gradually become modified by soil organisms. Plant roots predominantly follow the irregular voids between the aggregates. With increasing bulk density and growth of crops their impact increases. Mesofauna enlarge necks between void systems or locally ingest fine-grained soil material. The macrofauna produce channel-like tracks on the surface of the aggregates and/or produce channel systems in aggregates, whether or not connected to the surface. The largest part of the voids occurring in arable topsoils are voids due to tillage (Figure 13). In topsoils of a conventional farming system in The Netherlands tillage voids during the years occupy about 90% of the total macroporosity




Figure 12. Settlement of soil aggregates after tillage. a: soil structure about a month after ploughing; b. soil structure after seedbed preparation (magnification a, b: x 2).

(> 30 μm diameter voids); in an integrated farming system they occupy 5-10% less as result of the increased impact of soil organisms (Boersma and Kooistra, 1994). Shrinkage cracks, starting from tillage voids, are included in this group, but they contribute only a minority of the volume occupied.

The equipment used in agricultural engineering has increased in size and weight over the last decades. This development has led to increased soil compaction and deformation of the groundmass and to rut formation. Existing voids are reduced in size and changed in shape during passage of wheels. They may become disrupted, closed or can disappear completely. A reduction of the volume occupied by voids with diameters > 100 μm by 3% (v/v) or more, and an increase on the volume of the voids < 100 μm is also common, resulting in an only slightly decreased total porosity (Kooistra, 1987). The stress exerted decreases with depth and continues below the tilled layer (Koolen et al., 1992). Maximum compaction, however, generally occurs between 4 to 8 cm depth




Figure 13. Proportion of biological induced voids as part of the total macroporosity (voids with diameters > 30 μm) in 1985, 1987 and 1990 in a conventional and integrated farming system.

(Tanaka et al., 1988), due to the deloading effect at the surface. This often results in the soil in the formation of sets of small horizontal cracks. These cracks can also be formed below the wheels driving over the furrow bottom during ploughing, which gives the platy appearance to ploughpans. Where these sets of cracks occur in the tilled zone most tillage voids have disappeared (Słowińska-Jurkiewicz and Domżał, 1991). When the soil is brought under

cultivation, the surface soil is regularly tilled and the land is bare during several months of the year. When tilled and bare, the impact of wind and rain often lead to a disintegration of soil coherence. This starts with the aggregates at the surface. The disintegrated soil material can form crusts at the surface. When transport takes place, the soil material can move along the surface as soil erosion or flow downward into the soil as internal slaking. Microscopic studies reveal that there are many types of erosional and depositional crusts occurring in all climates (Courty, 1986; Kooistra and Siderius, 1986; Valentin and Ruiz Figueroa, 1987; Valentin and Bresson, 1992). Crusts cause restricted infiltration and limited plant germination due to their sorted, often laminated composition, lacking continuous voids. Transport of soil material along the surface due to wind and water erosion occurs on a large scale and is one of the main processes that takes place in areas with increased desertification. Also internal slaking occurs on a large scale. In the American soil classification "Soil Taxonomy" (Soil Survey Staff, 1975) a special diagnostic horizon, the agric horizon, has been defined to characterize the results of internal slaking. The illuviated soil material from the surface is deposited in voids where it forms coatings or complete infillings. The illuviated material can be deposited at any depth in the profile, depending on the present voids, quantities of water and soil material, and kind of soil material. Internal slaking results in a decrease of the total porosity and reduction of the continuous voids, which are essential for maintenance of water and air regimes and transport processes in the soil.

Repeated tillage also causes fragmentation, deformation and compaction of features due to the human impact. Although meant to loosen and equalize the soil, the aggregates formed by tillage can contain or consist of surface crusts, compacted layers, as well as soil material with infilled voids due to internal slaking. Moreover the distribution of crop residues etc. is also not evenly distributed. The resulting soil material is therefore often far more heterogeneous than natural soils.

5. Preferential Flow Patterns

In soils with large non-capillary sized voids, water and solutes flow via these voids. Large voids are cracks or planes formed by physical aggregate formation of soil tillage operations, or channels produced by roots or soil macrofauna. Coloured dyes and other traces are used by many investigators to determine and analyze large water-conducting voids (e.g. Ehlers, 1975; Bouma et al., 1977; Kooistra et al., 1985) The continuous water-conducting voids usually constitute only a few percent of the total void space and void necks have a substantial effect on the rate of flow. Dead-end voids and void necks are important locations where water with solutes can infiltrate into the soil matrix. Depending on the continuity of the macrovoids, water can bypass large volumes of soil

(e.g. Bouma and Dekker, 1978; Bouma et al., 1982; Van Stiphout et al., 1987). Due to this bypass flow the effect of added fertilizers or pesticides can be restricted as their solutes can be leached deep into the soil (e.g. Dekker and Bouma, 1984; Steenhuis et al., 1990) and can cause contamination of groundwater.

Preferential flow of water in unsaturated soil need not to be restricted to large void systems. Also in zones without large void systems, the water and solutes often infiltrate in fingered patterns. Vegetation can cause uneven infiltration of rain water. Isolated trees with a 'funnel' shaped canopy, which have a high rate of stem flow, can cause deep water infiltration under their stem as found in African savanna vegetation (Knapp, 1973). These wetting patterns can, besides vegetation, also be caused by microtopography, soil layering and water repellency. When fingered flow occurs, the spatial variability in soil moisture content can be very high (e.g. Dekker and Ritsema, 1995). These patterns can be found in sandy soils as well as in clay and peat soils.

Also on a smaller scale preferential water flows have various consequences. Depending on the local soil architecture, increasingly smaller voids receive water that form films along walls or fill small voids. In these environments, microbial activities take place, depending on the presence of substrate, water tension and related diffusion of solutes and gasses, and the distribution of void diameters. McGill and Myers (1987) calculated that only about 40% of the volume of voids is small enough to retain water and is large enough to permit entry of microbes. Combined with the above mentioned decisive factors, microbial activity would take place in a smaller percentage of the occurring voids. Moreover, microorganisms are not equally sensitive to the water potential for their activities. Consequently, microenvironmental conditions are often so heterogeneous that in spots near to each other simultaneous apparently exclusive processes can take place, e.g. N mineralization or immobilization. In general, the presence and accessibility (which implies the distribution of continuous void-systems) of substrate, and the preferential wetting patterns are the main factors determining microbial processes.

C. Processes Decreasing Heterogeneity

Processes leading to heterogeneity can, when occurring at higher intensities and/or over a longer time-span, lead to homogeneity as well. This may be true, for example for the impact of soil fauna and for human impact by cultivation practices. A reduction of heterogeneity can be due to the fact that a process, such as channel formation, is gradually occurring in the whole soil matrix. A process which directly leads to a decrease of heterogeneity is diffusion of chemical compounds and organisms in the soil, which is driven by concentration gradients.

1. Impact of Soil Fauna

Activities of mobile as well as sessile soil fauna can reduce soil heterogeneity. Mobile soil fauna in search for food move through the soil following existing void systems, enlarging pore necks where the passage is hampered and/or creating new channels. They exploit local zones where food is available and produce excrements on their way. In arable land, food sources are related to the root systems of the crops, incorporated organic matter and organic matter deposited at the surface. As soil organisms are generally strongly related to specific zones where the same moisture and temperature conditions prevail, the same architecture can develop in whole zones. A characteristic example is a moder humus zone in a spodosol (Figure 14a), in The Netherlands, which often occurs below a plaggen epipedon. The organic matter together with the fine-grained mineral material becomes conversed into shaped organo-mineral excrements produced by small soil fauna, chiefly microarthropods such as enchytraeids and collembola. The excrements occur in open or close packed, individual or slightly welded clusters between the larger mineral grains. Nearly all fine-grained mineral material is present in the organo-mineral excrements.

Sessile soil fauna create their own channel system which is in use for prolonged periods. They, however, also move around in search for food, but retreat into their own channels. These animals deposit their excreta at the surface or, less often, into old side-branches of their channel system. With time and presence of enough food homogeneous zones are formed. A characteristic example of this type is a mull humus in a mollic epipedon (Figure 14b). In a mull humus the organic material is intimately mixed with the mineral material forming a clay-humus complex. This type of humus form is produced by earthworms, who consume organic as well as mineral material. The excreta form clusters or small aggregates of strongly welded or shapeless material.

2. Impact of Diffusion

Diffusion is driven by, but also reduces concentration gradients. Diffusion is thus one of the major processes leading to homogeneity of the soil, although complete uniformity is approached at an infinitely slow rate, and new disruptions will have taken place before the soil is completely uniform. To get a feeling for the speed at which such homogenization occurs, Figure 15 shows the time course of the average concentration of a sphere of soil of 1 cm radius, inside a second sphere of 20 cm radius (see appendix for equations). At time zero, the concentration in the inner sphere is 1.0 and in the surrounding sphere 0. When we take typical values for the diffusion coefficient representing nitrate in a soil at field capacity, diffusion is so rapid, that the soil will be uniform for all




Figure 14. Humus forms produced by soil fauna. a. black moder humus; b. mull humus (magnification a: x 62; b: x 25).

practical purposes within a few days. For K it may be one or a few weeks, but for phosphate it will take weeks or months, depending on the P adsorption constant of the soil to reduce the concentration in the inner sphere to 50% of its original value. The time is scaled with the effective diffusion constant, and a tenfold reduction in mobility, leads to a tenfold increase in the time required for a set degree of homogenization. In drier soil conditions, the mobility of nutrients is reduced. So and Nye (1989) showed that for a tenfold decrease in D* from its value at field capacity (pF = 2.0) a sandy loam has to dry out until pF 3.3




Figure 15. Concentration decrease by diffusion from a 'hot spot' (initial concentration 1), consisting of an inner sphere of radius 1 cm in a 'cold' sphere (initial concentration 0) of radius 20 cm, with values of the effective diffusion constant 1, 0.1 and 0.01-0.001 $cm^2\ day^{-1}$, taken to be representative for nitrate, potassium and phosphate, respectively. Calculations based on equation (3) in the appendix.

and a silty clay until pF 4.5. Such a decrease in soil water content renders the diffusion of NO_3 in a dry soil similar to that of K at field capacity.

Homogenization of concentration by diffusion also depends on the absolute size of the 'hot spot'; in the spherical model the time required for homogenization is (approximately) proportional to the second power of the radius. Hot spots of nitrate of 3.2 cm radius thus will follow the line indicated for K hot spots of 1 cm radius. Heterogeneities at short distances are thus of practical relevance for solutes with an effective mobility as low as, or lower than that of inorganic phosphate. The diffusivity of oxygen in water is 10^4 times slower than that in free air, but the diffusivity of oxygen in a water saturated soil is still similar to

that of nitrate. Equilibration of oxygen concentration in a soil with aggregate sizes of 1 to 3 cm radius is thus a matter of hours to days (weeks), at least in the absence of oxygen consumption. With a continuing oxygen demand local differences in oxygen concentration and even local anaerobiosis can persist in (nearly) water saturated soil. The overriding importance of consumption on maintaining steep gradients was also shown for glucose diffusion into soil aggregates by Priesack and Kisser-Priesack (1993).

IV. Consequences of Heterogeneity for Processes and Resulting Patterns at Higher Levels of Complexity

This section is focused on the upper part of the chain as given in Figure 7. To what extent does soil structure and the distribution of soil carbon influence processes at a higher level of organization, such as water and nutrient use efficiency of agricultural systems? If a 'good' soil structure is to help in reducing negative environmental impact of agriculture, how does this work and how can it be used to the full extent?

Water and nutrient use efficiency of an agricultural system primarily depends on how well supply and demand are mutually adjusted. As long as demand exceeds supply, resource use efficiencies are generally high. It is difficult to achieve a high efficiency with a resource supply which does not limit productivity. Yet, highly productive but at the same time efficient agricultural production systems are needed for the next century. The 'efficiency gap' between well maintained experimental plots and current farmer's practice (Whitmore and Van Noordwijk, 1995) is at least partly due to difficulties in predicting rates of biologically mediated soil processes and to field scale spatial variability beyond the area exploited by an individual plant. Several of the components of the demand-supply balance are related to soil structure: root growth, mineralization, denitrification and leaching all are influenced by soil structure. To close the 'efficiency gap' a better quantitative understanding of the underlying patterns and processes is needed. In this context we may come back to the question of whether models can be based on the concept of a perfectly mixed homogeneous medium responding uniformly to outside influences or should be based on the recognition of soil as a highly structured, heterogeneous environment with large zones of little activity and scattered 'hot spots' full of activity. At first sight, the apparent success of many model descriptions of major soil processes, based on a homogeneous soil concept seems to indicate that the heterogeneity is more of academic than of practical interest.

The majority of transport models consider soil heterogeneity implicitly rather than explicitly. Relevant parameters are normally measured on the scale of an 'elementary soil volume' which can be thought to be representative. Thus, a

soil may be described as if it were homogeneous, but with an average gas diffusivity assigned to it which is based on the macropores, a water diffusivity dominated by the mesopores, and a by-pass flow description, based on the continuous macro-voids. The way in which these overall parameters are influenced by soil structure, and by variations in soil structure due to management actions remains obscure in many instances.

Models of nutrient uptake are normally based on assumptions of a regular root distribution. The consequences for uptake processes of non-regular root distribution and incomplete root-soil contact imposed on the root architecture by a structured soil can now be quantified (Van Noordwijk et al., 1993a; Veen et al., 1992). To a considerable extent, such effects of soil structure can be expressed as a reduction factor on the measured total root length and can be overcome by the plant by making more roots. Such reduction factors can give an alternative explanation to the reduction factors employed to estimate the 'live' fraction of a root system, but the two aspects are as yet difficult to separate and quantify. In many cases, models are used to evaluate the consequences of root length densities actually found in the soil and do not attempt to predict how root development depends on soil conditions, as mediated by soil structure.

Mineralization patterns do depend on soil structure. The consequences for food web functioning of the restricted accessibility of bacteria in small sized pores for protozoan predators are a recent focus of attention (Hassink et al. 1993; Wright et al. 1993). Soil food web models (De Ruiter et al., 1994) so far are remarkably similar to those for aquatic systems, and do not explicitly consider spatial structure of a soil, and the 'hot spot' nature of organic substrates. With the existing food web models, the measured N mineralization patterns in field soils can be predicted within a factor-two error range and they thus form an alternative to the C pool based model approach. Again, their relative success does not implicate that soil heterogeneity is not important for the functioning of the soil food web, as the calculations are based on actually observed population densities and do not attempt to predict the population development as such. Some mineralization models based on C pool sizes do include the concepts of 'physical protection' of soil carbon against bacteria and of bacteria against grazers (Verberne et al., 1990) but the protection factors are not yet completely operationally defined. Present attempts to do so are largely confined to soil texture and do not yet attempt to include soil structure. In the next generation of models, the mobility of soil organisms and their relations with soil structure may be more fully described.

Leaching rates critically depend on 'by-pass flow' through macropores. Considerable progress in field measurement of these properties has been made (Edwards et al., 1993). Despite considerable progress at the single aggregate level (Leffelaar, 1993), our understanding of denitrification, and its relation with measurable aspects of soil architecture is still far from comprehensive. Development of anaerobic sites and denitrification depend on the presence of

decomposable organic substrates under conditions of slow oxygen replenishment. Soil structure and soil water content determine the oxygen transport rates, but the spatial distribution of oxygen demanding bacteria is probably related to the hot spots of organic substrates.

Using larger amounts of organic inputs is not a simple recipe to obtain a better soil structure and a more efficient and yet productive type of agriculture. If they are integrated with other aspects of crop management, however, biological means of maintaining and improving soil structure can give a real contribution to overall resource use efficiency. Biological management of soil structure by earthworm and root channels remains an exciting option (Dexter, pers. comm.) for extensive land use systems; one should not expect miracles however and where the soil building has been severely mistreated, e.g. during deforestation in the tropics, the biological soil architects will take a very long time to reconstruct anything similar to the original structure.

V. Importance of Heterogeneity for Carbon Retention in Soil

The organic carbon present in the soil is derived from aboveground organic materials incorporated as part of tillage activities or below ground organic inputs from primary producers (see part I and Figure 2). The incorporated organic inputs have a heterogeneous distribution. The method of incorporation determines the spatial pattern: e.g. inverted soil surfaces deposited by plough shares (Figure 3), more finely mixed material with other tillage implements. The same is true for the below ground organic inputs from primary producers, viz. leaves pulled into wormholes or roots with a distinct distribution pattern in the soil, whether or not redistributed by tillage activities. This heterogeneous distribution by itself may not be very important for decomposition rates or its complement, carbon retention in soil. The same organic material can be decomposed via different pathways and in different time lapses. Thus it is not the location as such of organic carbon which matters, but its effects on accessibility of the organic input for decomposing bacteria and fungi, predators and water, gasses and solutes. If the accessibility is affected, e.g. because a crop residue is incorporated in groundmass, decomposition can be very slow, while residues a few cm away along a crack are quickly fragmented and partly consumed (Figure 16). The accessibility for incorporated organic matter differs strongly and cannot be predicted. Those of the original below ground organic materials, such as roots, exudates and mycorrhizal hyphae are in principle far more accessible, as their locations are determined by both the root distribution pattern itself and the root preference for existing void systems.

In more natural arable systems (e.g. in agroforestry), faunal activity is the major determinant of incorporation of above ground organic inputs. Considerable differences in litter layer exist between vegetation types, but also over small




Figure 16. Difference in accessibility of crop residues in the plowlayer; the crop residue along the crack is partly consumed and fragmented by enchytraeids; the residue embedded in the groundmass remains intact (magnification x 2).

distances within a single forest type. Burghouts (1993) analyzed such patterns for a Bornean rainforest and concluded that differences in deposition rates rather than turnover rates were responsible for the apparent differences in carbon retention on the soil surface. Within a structured soil, however, possibilities for differences in turnover rates are substantially larger than on the soil surface.

Primary organic matter is decomposed faster than secondary organic matter present in excreta. The most decomposable excrements are those of the large macrofauna such as earthworm excreta. Mesofauna (enchytraeids, mites and collembola) consume coarser fragments of the incorporated organic material, sometimes leaving their excrements in the cavity they produced. Excrements can be conserved in the soils for centuries, in spite of tillage activities, without microscopic recognizable changes (e.g. Kooistra, 1978). Otherwise excrements disintegrate slowly forming shapeless organic coatings and bridges between mineral grains as in moder podsol soils.

VI. Summary and Conclusions

Soil structure has a biological component, the purposely made architectural part, which is highly variable in quality and quantity, occurs at different scales and varies throughout the year. The biological impact of belowground communities and plant roots depends on the different requirements and can be mutually exclusive, mutually complementary or dependent, completely independent or mixtures of these. The biological component of soil structure, therefore, is far

less predictable than the physical part of soil structure dealing with swell/shrink and freeze/thaw cycles, and the human impact by tillage and management operations.

Nearly every observable/measurable biotically related soil phenomenon has a heterogeneous distribution in the soil. Increase and decrease of heterogeneity can be based on similar processes, but at different time scales, or on different processes. Recognition of and understanding of the dynamics of these heterogeneities appears to be essential for a 'real' understanding of what's going on in the soil. Yet, for many purposes models which treat soils as (a number of) black boxes, with average rates assigned to them have been highly successful. Such models are based on empirical 'fudge' factors, which are not always properly acknowledged in so-called 'process-based' models. Progress will be made when the empirically determined average parameter values for the black boxes can be explained on the basis of the underlying heterogeneity, the real actors and the variation in internal heterogeneity between sites. A number of new concepts and methods is now available for such an excercise.

The heterogeneous distribution by itself is not the main controlling factor in regard to decomposition rates or carbon retention in soils. Rather, it is the accessibility (not the location) of the organic input for decomposing bacteria and fungi, predators, water, gasses and solutes which is of primary importance. In studying these phenomena, new methods for analyzing and interpreting distribution patterns of single features across a range of scales are needed, especially methods to quantify the spatial correlations of patterns and their significance for turnover of organic inputs in the soil. As some of the products of turnover (e.g. anaerobically formed gasses such as CH_4 and N_2O, and solutes such as NO_3) are definitely undesirable when they reach other compartments of the environment (e.g. atmosphere, groundwater), research on soil structure may gain new relevance.

References

Altemüller, H.-J. and Th. Haag. 1983. Microscopische Untersuchungen an Maiswurzeln im umgestorten Bodenverband. *Kali-Briefe* 16: 349-363.

Altemüller, H.-J. and B. Van Vliet-Lanoë. 1990. Soil thin section fluorescence microscopy. p. 565-581. In: L.A. Douglas (ed.), *Soil Micromorphology*: a Basic and Applied Science. Developments in Soil Science, 19. Elsevier, Amsterdam.

Anderson, J.M. and J.S.I. Ingram (eds.), 1993. *Tropical soil biology and fertility, a Handbook of Methods*. 2nd edition. CAB International, Wallingford, 221 pp.

Beare, M.H. and R.R. Bruce. 1993. A comparison of methods for measuring water-stable aggregates: implications for determining environmental effects on soil structure. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma*, 56: 87-104.

Bennett, R.H., W.R. Bryant, and M.H. Hulbert (eds.), 1990. *Microstructure of Fine-Grained Sediments*. Springer, New York, U.S.A., 582 pp.

Blokhuis, W.A., M.J. Kooistra, and L.P. Wilding. 1990. Micromorphology of cracking clayey soils (Vertisols). p. 123-148. In: L.A. Douglas (ed.), *Soil Micromorphology: a Basic and Applied Science*. Developments in Soil Science, 19. Elsevier, Amsterdam.

Bodemkaart. 1979. *Bodemkaart van Nederland, schaal 1:50 000*. Blad 34 W/O met toelichting. Stiboka, Wageningen, The Netherlands, 166 pp.

Boersma, O.H. and M.J. Kooistra. 1994. Influences on soil structure of silt loam typic fluvaquents under various agricultural management practices. *Agric. Ecosyst. Environ.* 51:21-42.

Böhm, W. 1979. *Methods of studying root systems*. Springer, Berlin, 188 pp.

Bouma, J., A. Jongerius, O.H. Boersma, A. Jager and D. Schoonderbeek. 1977. The function of different types of macropores during saturated flow through 4 swelling soil horizons. *Soil Sci. Soc. Am. J.* 41 (5): 945-950.

Bouma, J. and L.W. Dekker. 1978. A case study on infiltration into dry clay soil. I. Morphological observations. *Geoderma* 20: 27-40.

Bouma, J. A. Jongerius and D. Schoonderbeek. 1979. Calculation of saturated hydraulic conductivity of some pedal clay soils using micromorphometric data. *Soil Sci. Soc. Am. J.* 43 (2): 261-264.

Bouma, J., C.F.M. Belmans and L.W. Dekker. 1982. Water infiltration and redistribution in a silt loam subsoil with vertical worm channels. *Soil. Sci. Soc. Am. J.* 46: 917-921.

Brussaard, L. and L.T. Runia. 1984. Recent and ancient traces of scarab beetle activity in sandy soils of The Netherlands. *Geoderma*. 34: 229-250.

Bullock, P., N. Federoff, A. Jongerius, G. Stoops, and T. Tursina. 1985. *Handbook for soil thin section description*. Waine Research Publ., Wolverhampton, U.K.

Burghouts, T. 1993. *Spatial heterogeneity of nutrient cycling in Bornean rainforest*. Free University Amsterdam, PhD Thesis, 156 pp.

Cambardella, C.A. and E.T. Elliott. 1993. Methods for physical separation and characterization of soil organic matter fractions. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma* 56: 449-457.

Carslaw, H.S. and J.C. Jaeger. 1959. *Conduction of heat in solids*. 2nd ed. Clarendon, Oxford.

Courty, M.A. 1986. Morphology and genesis of soil surface crusts in semi-arid conditions (Hissar region, Northwest India). p. 32-40. In: Callebaut, F., D. Gabriels, and M. De Boodt (eds.), *Assessment of soil surface sealing and crusting*. State Univ. Ghent, Belgium.

Dekker, L.W. and J. Bouma. 1984. Nitrogen leaching during sprinkler irrigation of a Dutch clay soil. *Agric. Water Manage.* 9: 37-45.

Dekker, L.W. and C.J. Ritsema. 1995. Fingerlike wetting patterns in two water repellent loam soils. *J. Env. Qual.* (in press).

De Ruiter, P.C., J.C. Moore, K.B. Zwart, L.A. Bouwman, J. Hassink, J. Bloem, J.A. de Vos, J.C.Y. Marinissen, W.A.M. Didden, G. Lebbink and L. Brussaard. 1994. Simulation of nitrogen mineralization in the belowground food webs of two winter wheat fields. *J. Appl. Ecol.* 30: 95-106.

De Willigen, P. and Van Noordwijk, M., 1987. *Roots, plant production and nutrient use efficiency*. Ph.D. Thesis, Agric. Univ. Wageningen, The Netherlands, 282 pp.

Dexter, A.R. 1988. Advances in characterization of soil structure. *Soil Tillage Res.* 11: 199-239.

Diggle, P.J. 1983. *Statistical Analysis of Spatial point Patterns*. Academic Press, London. 148 pp.

Edwards, W.M., M.J. Shipitalo, and L.B. Owens. 1993. Gas, water and solute transport in soils containing macropores: a review of methodology. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma*, 57: 31-49.

Ehlers, W. 1975. Observations on earthworm channels and infiltration on tilled and untilled loess soils. *Soil Sci.* 119: 242-249.

Elliott, E.T. 1986. Aggregate structure and carbon, nitrogen and phosphorus in native and cultivated soils. *Soil Sci. Soc. Am. J.* 50: 627-633.

Gupta, V.V.S.R. and J.J. Germina. 1988. Distribution of microbial biomass and its activity in different soil aggregate classes as affected by cultivation. *Soil Biol. Biochem.* 2 (6): 777- 786.

Hassink J., L.A. Bouwman, K.B. Zwart, J. Bloem, and L. Brussaard. 1993. Relationships between soil texture, physical protection of organic matter, soil biota, and C and N mineralization in grassland soils. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma* 57: 105-128.

Hassink, J. 1995. Density fractions of soil macroorganic matter and microbial biomass as predictors of C and N mineralization. *Soil Biol. Biochem* (in press).

Jongerius, A. and G. Heintzberger. 1975. *Methods in soil micromorphology; a technique for the preparation of large thin sections*. Soil Survey Paper no. 10, Netherlands Soil Survey Institute, Wageningen, 48 pp.

Kemper, W.D. and E.J. Koch. 1966. *Aggregate stability of soils from Western United States and Canada*. Techn. Bull., 1355, Agricultural Research Service, U.S. Dept. of Agriculture, Washington, D.C., 52 pp.

Knapp, R. 1973. *Die Vegetation von Africa*. Gustav Fischer Verlag, Stuttgart, 626 pp.

Kooistra, M.J. 1978. *Soil development in recent marine sediments of the intertidal zone in the Oosterschelde, The Netherlands. A soil micromorphological approach*. Soil Survey Paper no. 14, Netherlands Soil Survey Institute, Wageningen, 183 pp.

Kooistra, M.J. 1982. Micromorphology. p. 71-89. In: Murthy, R.S., L.R. Hirekerur, S.B. Deshpande, and B.V. Venkata Rao (eds.), *Benchmark Soils of India. Morphology, Characteristics and Classification for Resource Management*. National Bureau of Soil Survey and Land Use Planning (ICAR) in collaboration with The All India Soil and Land Use Survey (Government of India). Agricultural Universities and Soil Survey Organizations of the State Departments of Agriculture.

Kooistra, M.J. 1987. The effects of compaction and deep tillage on soil structure in a Dutch sandy loam soil. p. 445-450. In: Fedoroff, N., L.M. Bresson, and M.A. Courly (eds.), *Soil Micromorphology*, AFES, Plaisir, France.

Kooistra, M.J. 1991. A micromorphological approach to the interactions between soil structure and soil biota. *Agric. Ecosyst. Environ.* 34: 315-328.

Kooistra, M.J., J. Bouma, O.H. Boersma, and A. Jager. 1985. Soil-structure differences and associated physical properties of some loamy Typic Fluvaquents in The Netherlands. *Geoderma* 36: 215-228.

Kooistra, M.J. and W. Siderius. 1986. Micromorphological aspects of crust formation in Savanna climate under rainfed subsistence agriculture. p. 9-17. In: Callebaut, F., D. Gabriels, and M. De Boodt (eds.), *Assessment of soil surface sealing and crusting*. State Univ. Ghent, Belgium.

Kooistra, M.J., G. Lebbink, and L. Brussaard. 1989. The Dutch Programme on Soil Ecology of Arable Farming Systems II. Geogenesis, agricultural history, field site characteristics and present farming systems at the Lovinkhoeve experimental farm. *Agric. Ecosyst. Environ.* 27: 361-387.

Kooistra, M.J., A.S.R. Juo, and D. Schoonderbeek. 1990. Soil degradation in cultivated alfisols under different management systems in southwestern Nigeria. p. 61-70. In: L.A. Douglas (ed.), *Soil Micromorphology: a Basic and Applied Science*. Developments in Soil Science, 19. Elsevier, Amsterdam.

Kooistra, M.J., D. Schoonderbeek, F.R. Boone, and M. Van Noordwijk. 1992. Root-soil contact of maize, as measured by thin-section technique. II. Effects of soil compaction. *Plant Soil* 139: 119-129.

Kooistra, M.J. and N.K. Tovey. 1994. Effects of compaction on soil microstructure. p. 91-111 In: Soane, B.D. and Van Ouwerkerk, C. (eds.), *Soil compaction in crop production.* Elsevier, Amsterdam, The Netherlands.

Koolen, A.J., P. Lerink, D.A.G. Kurstjens, J.J.H. Van den Akker, and W.B.M. Arts. 1992. Prediction of aspects of soil-wheel systems. *Soil Till. Res.* 24:381-396.

Kuipers, H., and C. Van Ouwerkerk. 1963. Total pore-space estimationsin freshly ploughed soil. *Neth. J. Agric. Sci.* 11, 1: 45-53.

Leffelaar, P.A., 1993. Water movement, oxygen supply and biological processes on the aggregate scale. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma* 57: 143-165

McGill, W.B. and R.J.K. Myers. 1987. Controls on dynamics of soil and fertilizer nitrogen. p. 74-99. In: R.F. Follett, J.W.B. Stewart, and C.V. Cole (eds.), *Soil fertility and organic matter as critical components of production systems.* SSSA Spec. Pub. no. 19.

Murphy, C.P., 1986. *Thin section preparation of soils and sediments.* A.B. Academic Publishers, Berkhamsted, U.K., 149 pp.

Meijboom, F.W., J. Hassink and M. Van Noordwijk. 1995. Density fractionation of soil macroorganic matter using silica suspensions. *Soil Biol. Biochem* (in press).

North, P.F., 1976. Towards an absolute measurement of soil structural stability using ultrasound. *J. Soil Sci.* 27: 541-459.

Pielou, E.C., 1969. *An Introduction to Mathematical Ecology,* Wiley-Interscience, New York. 286 pp.

Postma, J., and H.-J. Altemüller. 1990. Bacteria in thin soil sections stained with the fluorescent brightener calcofluor white M2R. *Soil Biol. Biochem.* 22:89-96.

Pons, L.J., and I.S. Zonneveld. 1965. *Soil ripening and soil classification. Initial soil formation in alluvial deposits and a classification of the resulting soils.* Int. Inst. Land Reclamation and Improvement, Publ. 13, Wageningen, The Netherlands, 128 pp.

Priesack E., and G.M. Kisser-Priesack. 1993. Modelling diffusion and microbial uptake of 13C-glucose in soil aggregates. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma* 56:561-573.

Słowińska-Jurkiewicz, A. and H. Domżał. 1991. The structure of the cultivated horizon of soil compacted by the wheels of agricultural ractors. *Soil Till. Res.* 19:215-226.

Smeerdijk, D.G., T. Spek, and M.J. Kooistra. 1994. Anthropogenic soil formation and agricultural history of the open fields of Valthe (Drenthe, The Nethertlands) in medieval and early modern times. Integration of palaeooecology, historical geography and soilscience. *Mededelingen Rijks Geologische Dienst.* (in press).

So, H.B., and P.H. Nye. 1989. The effect of bulk density, water content and soil type on the diffusion of chloride in soil. *J. Soil Sci.* 40: 743-749.

Soil Survey Staff. 1975. Soil Taxonomy. *A basic system of soil classification for making and interpreting soil surveys.* US Government Printing Office, Washington, D.C., U.S.A., 754 pp.

Staricka, J.A., R.R. Allmaras, and W.W. Nelson. 1991. Spatial variation of crop residue incorporated by tillage. *Soil Sci. Soc. Am. J.* 55:1668-1674.

Steenhuis, T.S., W. Staubitz, M.S. Andreini, J. Surface,T.L. Richard, R. Paulsen, N.B. Pickering, J.R. Hagerman and L.D. Geohring. 1990. Preferential movement of pesticides and tracers in agricultural soils. *J. Irrig. Drain. Eng.* 116:50-66.

Tanaka, T., M. Yamazaki, and A.T. Abebe. 1988. Multiple pass effect of a wheel on soil compactability. In: Gee-Glough, D. and V.M. Salokke (eds.), *Proceedings of 2nd Asian- pacific Conference of the International Society for Terrain-vehicle Systems.* Asian Institute of Technology, Bangkok.

Tipkötter, R., K. Ritz and J.F. Darbyshire. 1986. The preparation of soil thin sections for biological studies. *J. Soil Sci.* 37:681-690.

Tisdall, J.M., and J.M. Oades. 1979. Stabilization of soil aggregates by the root systems of rye-grass. *Aust. J. Soil Res.* 17: 429-441.

Valentin, C. and J.F. Ruiz Figueroa. 1987. Effects of kinetic energy and water application rate on the development of crusts in a fine sandy loam soil using sprinkling irrigation and rainfall simulation. p. 401-409. In: Fedoroff, N., L.M. Bresson, and M.A. Courty (eds.), *Soil Micromorphology,* AFES, Plaisir, France.

Valentin, C. and L.M. Bresson. 1992. Morphology, genesis and classification of surface crusts in loamy and sandy soils. *Geoderma* 55: 225-246.

Van Noordwijk M., Widianto, M. Heinen, and K. Hairiah. 1991. Old tree root channels in acid soils in the humid tropics: important for crop root penetration, water infiltration and nitrogen management. *Plant Soil* 134: 37-44.

Van Noordwijk, M., M.J. Kooistra, F.R. Boone, B.W. Veen, and D. Schoonderbeek. 1992. Root-soil contact of maize, as measured by thin section technique. I. Validity of the method. *Plant Soil* 139: 109-118.

Van Noordwijk, M., G. Brouwer, K. Harmanny. 1993a. Concepts and methods for studying interactions of roots and soil structure. In: Brussaard, L. and M.J. Kooistra (eds), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma* 56: 351-375.

Van Noordwijk, M., G. Brouwer, P. Zandt, F.W. Meijboom, and S. Burgers. 1993b. *Root patterns in space and time: procedures and programs for quantification.* IB-Nota 268, IB-DLO Haren.

Van Noordwijk, M., P.C. De Ruiter, K.B. Zwart, J. Bloem, J.C. Moore, H.G. Van Faassen, and S. Burgers. 1993c. Synlocation of biological activity, roots, cracks and recent organic inputs in a sugar beet field. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma* 56: 265-276.

Van Noordwijk, M., D. Schoonderbeek, and M.J. Kooistra. 1993d. Root-soil contact of field grown winter wheat. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma* 56:227-287.

Van Stiphout, T.P.J., H.A.J. van Lanen, O.H. Boersma and J. Bouma. 1987. The effect of bypass flow and internal catchment of rain on the water regime in a clay loam grassland soil. *J. Hydrol.* 95, 1/2: 1-11.

Veen B.W., M. Van Noordwijk, P. De Willigen, F.R. Boone and M.J. Kooistra. 1992. Root-soil contact of maize, as measured by thin-section technique. III. Effects on shoot growth, nitrate and water uptake efficiency. *Plant Soil* 139: 31-138.

Verberne, E.L.J., J. Hassink, P. De Willigen, J.J.R. Groot, and J.A. Van Veen. 1990. Modelling organic matter dynamics in different soils. *Neth. J. Agric. Sci.* 38: 221-238.

Vittorello, V.A., C.C. Cerri, F. Andreux, C. Feller, and R.L. Victoria. 1989. Organic matter and natural carbon-13 distribution in deforested and cultivated oxisols. *Soil Sci. Soc. Am. J.* 53: 773-778.

Wright D.A., K. Killham, L.A. Glover, and J.I. Prosser. 1993. The effect of location in soil on protozoal grazing of a genetically modified bacterial inoculum. In: Brussaard, L. and M.J. Kooistra (eds.), Int. Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships. *Geoderma* 56: 633-640.

Whalley, W.B. (ed.), 1979. Scanning Electron Microscopy in the Study of Sediments. *Geoabstracts*, Norwich, U.K., 414 pp.

Whitmore, A.P.M. and M. Van Noordwijk. 1995. Bridging the gap between environmentally acceptable and economically desirable nutrient supply. *Proceedings Long Ashton Symposium.* Sept. 1993. (in press).